

AMENDMENTS TO THE CLAIMS

- 1) (currently amended) A process to recover carotene concentrates ~~concentrates~~ comprising the steps of:
 - i) subjecting alkyl esters produced from palm oil to at least one stage vacuum distillation at temperature ranging from 80-220°C and pressure at less than 40mTorr to yield phytonutrients concentrate in residue;
 - ii) separation of polar lipids and other impurities from the residue in step (i);
 - iii) subjecting the treated residue from step (ii) to a second vacuum distillation wherein residue from the distillation contain carotenes concentrates consisting carotenes; ubiquinones and phospholipids₇.
- 2) (original) A process to recover carotene concentrates claimed in claim 1 wherein the second vacuum distillation is carried out at temperature ranging from 80°C to 200°C and at pressure less than 40mTorr.
- 3) (original) A process to recover carotene concentrates as claimed in claim 1 wherein the separation of polar lipids and other impurities in step (ii) is done in any one of the ways consisting of:
 - i) treating the residue in step (i) of claim 1 with a hydrocarbon solvent with or without subsequent alkyl alcohol/treated water purification to remove the monoacylglycerols; or
 - ii) re-transesterifying the residue in step (i) of claim 1 using alkaline catalysts to convert the traces of acylglycerols into alkyl esters and glycerol; or
 - iii) direct filtration of residue in step (i) of claim 1 under vacuum.

- 4) (original) A process to recover carotene concentrates as claimed in claim 3 wherein the mixture in step (i) is chilled down to low temperature for at least 2 hours and monoacylglycerols is separated from the residue.
- 5) (original) A process to recover carotene concentrates as claimed in claim 3 wherein alkaline catalyst used in the re-transesterification in step (ii) is selected from a group consisting of such as sodium hydroxide, potassium hydroxide and sodium methylate in the presence of short and branched alkyl alcohol such as methanol and ethanol.
- 6) (original) A process to recover carotene concentrates as claimed in claim 3 wherein 2% of palm oil ethyl esters are added to the treated residue in step (iii) prior to subsequent vacuum distillation.
- 7) (original) A process to recover carotene concentrates as claimed in claim 1 wherein the carotenes concentrate is further purified by either:
 - i) adding alkaline catalysts in presence of alkyl alcohol such as potassium hydroxide in ethanol; or
 - ii) adding hydrocarbon solvent and alkyl alcohol and chilled to -10°C for at least one hour to partition the carotenes into hydrocarbon layer.
- 8) (currently amended) A process to recover carotene concentrates as claimed in claim 3 ~~or~~ and 7 wherein the hydrocarbon solvents used are hexane or iso-octane and the alkyl alcohols used are short and branched alkyl alcohols such as methanol and ethanol.
- 9) (original) A process to recover carotene concentrates as claimed in claim 1 wherein the alkyl esters is produced from the crude palm oil or treated palm oil such as bleached and

degummed palm oil and membrane filtered palm oil.

- 10) (original) A process to recover carotene concentrates as claimed in claim 9 wherein the removal of excess of alkaline catalyst in alkyl esters produced is carried out by using acidified water pH between 4-5.
- 11) (currently amended) Carotenes, ubiquinones, and phospholipids as produced in ~~any of the claims 1 to 10~~ claim 1.